



Datum: 22.02.2017
Kontakt: Ing. Veronika Iro, B.Sc.
Abteilung: REGA
Tel. / Fax: +43 (0) 505 55 – 36247
E-Mail: pv-implementation@ages.at
Unser Zeichen: PHV-9806001-A-170222
Ihr Zeichen:

PHV-issue: Ethinylestradiol

Sehr geehrte Damen und Herren,

In der Sitzung vom Januar 2017 kam das CMDh zu dem Schluss, europaweit die Fach- und Gebrauchsinformation aller Ethinylestradiol-hältigen kombinierten hormonalen Kontrazeptiva zu ändern. (siehe: <http://www.hma.eu/249.html>)



**CMDh RECOMMENDATION
COMBINED HORMONAL CONTRACEPTIVES (CHCs)
CONTAINING ETHINYLESTRADIOL**

Inclusion of a contra indication regarding an interaction between combined hormonal contraceptives containing ethinylestradiol with the medicinal products containing ombitasvir/paritaprevir/ritonavir (Viekirax) and dasabuvir with or without ribavirin agreed by CMDh in January 2017

SmPC

	SmPC for all CHCs containing ethinylestradiol
Section 4.3	<CHC trade name> is contraindicated for concomitant use with the medicinal products containing ombitasvir/paritaprevir/ritonavir and dasabuvir, (see sections 4.4 and section 4.5).
Section 4.4	<u>ALT elevations</u> During clinical trials with patients treated for hepatitis C virus infections (HCV) with the medicinal products containing ombitasvir/paritaprevir/ritonavir and dasabuvir with or without ribavirin, transaminase (ALT) elevations higher than 5 times the upper limit of normal (ULN) occurred significantly more frequent in women using ethinylestradiol-containing medications such as combined hormonal contraceptives (CHCs) (see sections 4.3 and 4.5).
Section 4.5	<u>Pharmacodynamic interactions</u> Concomitant use with the medicinal products containing ombitasvir/paritaprevir/ritonavir and dasabuvir, with or without ribavirin may increase the risk of ALT elevations (see sections 4.3 and 4.4). Therefore, <CHC trade name>-users must switch to an alternative method of contraception (e.g., progestagen-only contraception or non-hormonal methods) prior to starting therapy with this combination drug regimen. <CHC trade name> can be restarted 2 weeks following completion of treatment with this combination drug regimen.

Package leaflet

	SmPC for all CHCs containing ethinylestradiol
Do not use <CHC trade name>	Do not use <CHC trade name> if you have hepatitis C and are taking the medicinal products containing ombitasvir/paritaprevir/ritonavir and dasabuvir (see also in section Other medicines and <CHC trade name>).
Other medicines and <CHC trade name>	Do not use <CHC trade name> if you have Hepatitis C and are taking the medicinal products containing ombitasvir/paritaprevir/ritonavir and dasabuvir as this may cause increases in liver function blood test results (increase in ALT liver enzyme). Your doctor will prescribe another type of contraceptive prior to start of the treatment with these medicinal products. <CHC trade name> can be restarted approximately 2 weeks after completion of this treatment. See section “Do not use <CHC trade name>”.